

The Cardy-Verlinde Formula and Charged Topological AdS Black Holes

Donam Youm¹

ICTP, Strada Costiera 11, 34014 Trieste, Italy

Abstract

We consider the brane universe in the bulk background of the charged topological AdS black holes. The evolution of the brane universe is described by the Friedmann equations for a flat or an open FRW-universe containing radiation and stiff matter. We find that the temperature and entropy of the dual CFT are simply expressed in terms of the Hubble parameter and its time derivative, and the Friedmann equations coincide with thermodynamic formulas of the dual CFT at the moment when the brane crosses the black hole horizon. We obtain the generalized Cardy-Verlinde formula for the CFT with an R-charge, for any values of the curvature parameter k in the Friedmann equations.

May, 2001

¹E-mail: youmd@ictp.trieste.it

Recently, there has been growing interest in holographic bounds in cosmology [1, 2, 3, 4, 5, 6], after the initial work by Fischler and Susskind [7]. The holographic bound in its original form is found to be violated by the closed Friedmann-Robertson-Walker (FRW) universe [7] and the later works attempted to circumvent such a problem through various modifications. Later, Verlinde [8, 9] proposed an ingenious holographic bound for a radiation dominated closed FRW universe which remains valid throughout the cosmological evolution. The Verlinde's holographic bound unifies the Bekenstein bound [10] for an weakly self-gravitating universe and the Hubble bound [1, 2, 5, 6] for a strongly self-gravitating universe in an elegant way. (Cf. It is shown in Refs. [11, 12] that the Verlinde's entropy bound is equivalent to causal entropy bound, which is a covariant formulation of the Hubble entropy bound.) Furthermore, it is shown [8] that the Cardy formula [13] for a two-dimensional conformal field theory (CFT) can be generalized to an arbitrary spacetime dimensions and such generalized formula, called the Cardy-Verlinde formula, coincides with the Friedmann equation at the moment when the proposed cosmological bound is saturated. Such results were generalized to brane universes in various bulk backgrounds [14, 15, 16, 17, 18, 19, 20, 21, 22, 23]. The quantum effects to the Cardy-Verlinde formula were studied in Refs. [24, 25, 26]. In our previous work [27], we showed that the results of Refs. [8, 9] continue to hold even for an open and a flat radiation dominated FRW universes induced on a brane moving in the bulk background of the topological AdS-Schwarzschild black holes. In this note, we study a generalization to the brane universe in the bulk background of the charged topological AdS black holes. (The results in this paper can be straightforwardly generalized to the multi-charged topological AdS black holes in string theories.) The holographic dual theory corresponds to the CFT with an R-charge. We show that the temperature and entropy of the CFT are still simply expressed in terms of the Hubble parameter and its time derivative when the brane crosses the black hole horizon. We find that the Friedmann equations coincide with thermodynamic formulas of the CFT at the moment when the brane crosses the black hole horizon. We generalize the Cardy-Verlinde formula to include the contribution from the R-charge of the CFT, for any values of the curvature parameter k in the Friedmann equations.

The charged topological AdS black hole solution in $(n + 2)$ -dimensions has the following form [28, 29, 30, 31, 32, 33, 34]:

$$\begin{aligned}
ds_{n+2}^2 &= -h(a)dt^2 + \frac{1}{h(a)}da^2 + a^2\gamma_{ij}(x)dx^i dx^j, \\
h(a) &= k - \frac{w_{n+1}M}{a^{n-1}} + \frac{nw_{n+1}^2Q^2}{8(n-1)a^{2n-2}} + \frac{a^2}{L^2}, \\
\phi &= \frac{n}{4n-4} \frac{w_{n+1}Q}{a^{n-1}}, \quad \omega_{n+1} = \frac{16\pi G_{n+2}}{n\text{Vol}(M^n)},
\end{aligned} \tag{1}$$

where γ_{ij} is the horizon metric for a constant curvature manifold M^n with the volume $\text{Vol}(M^n) = \int d^n x \sqrt{\gamma}$, G_{n+2} is the $(n + 2)$ -dimensional Newton's constant, M is the ADM mass of the black hole, Q is the electric charge, ϕ is the electrostatic potential

(difference between the horizon and infinity), and L is the curvature radius of the background AdS spacetime. The horizon geometry of the black hole is elliptic, flat and hyperbolic for $k = 1, 0, -1$, respectively. The Bekenstein-Hawking entropy S , the chemical potential ϕ_H and the Hawking temperature $\mathcal{T}$ of the black hole are

$$\begin{aligned} S &= \frac{a_H^n \text{Vol}(M^n)}{4G_{n+2}}, & \phi_H &= \frac{n}{4n-4} \frac{w_{n+1}Q}{a_H^{n-1}}, \\ \mathcal{T} &= \frac{h'(a_H)}{4\pi} = \frac{(n+1)a_H^2 + (n-1)kL^2}{4\pi L^2 a_H} - \frac{nw_{n+1}^2 Q^2}{32\pi a_H^{2n-1}}, \end{aligned} \quad (2)$$

where a_H is the horizon, defined as the largest root of $h(a) = 0$, and the prime denotes the derivative w.r.t. a .

We obtain the effective Friedmann equations describing the evolution of the brane universe in the above bulk background (1). In terms of a new time coordinate τ , satisfying

$$\frac{1}{h(a)} \left(\frac{da}{d\tau} \right)^2 - h(a) \left(\frac{dt}{d\tau} \right)^2 = -1, \quad (3)$$

the induced metric of the n -brane takes the standard Robertson-Walker form

$$ds_{n+1}^2 = -d\tau^2 + a^2(\tau) \gamma_{ij} dx^i dx^j, \quad (4)$$

with the cosmic scale factor a . Making use of Eq. (3) and the following equation of motion [9] for the brane action:

$$\frac{dt}{d\tau} = \frac{\sigma a}{h(a)}, \quad (5)$$

where the parameter σ is related to the brane tension, we obtain the following Friedmann equation:

$$H^2 = \frac{\omega_{n+1}M}{a^{n+1}} - \frac{nw_{n+1}^2 Q^2}{8(n-1)a^{2n}} - \frac{k}{a^2}, \quad (6)$$

after setting $\sigma = 1/L$. Here, $H \equiv \dot{a}/a$ is the Hubble parameter, where the overdot denotes the derivative w.r.t. τ . Taking the τ -derivative of Eq. (6), we obtain the second Friedmann equation

$$\dot{H} = -\frac{n+1}{2} \frac{\omega_{n+1}M}{a^{n+1}} + \frac{n^2 w_{n+1}^2 Q^2}{8(n-1)a^{2n}} + \frac{k}{a^2}. \quad (7)$$

So, the motion in the bulk background (1) induces radiation matter $\sim M/a^{n+1}$ and stiff matter $\sim Q^2/a^{2n}$ [17] on the brane universe.

Making use of the fact that the metric for the boundary CFT can be determined only up to a conformal factor [35, 36], we rescale the boundary metric for the CFT to be of the following form:

$$ds_{CFT}^2 = \lim_{a \rightarrow \infty} \left[\frac{L^2}{a^2} ds_{n+2}^2 \right] = -dt^2 + L^2 \gamma_{ij} dx^i dx^j. \quad (8)$$

It is argued in Ref. [37] that thermodynamic quantities of the CFT at high temperature can be identified with those of the bulk AdS black hole. Note, the CFT time is rescaled by the factor L/a w.r.t. the AdS time. So, the energy E , the chemical potential Φ_H and the temperature T of the CFT are rescaled by the same factor L/a :

$$\begin{aligned} E &= M \frac{L}{a}, & \Phi_H &= \phi_H \frac{L}{a} = \frac{n}{4n-4} \frac{w_{n+1} Q L}{a_H^{n-1} a}, \\ T &= \mathcal{T} \frac{L}{a} = \frac{1}{4\pi a} \left[(n+1) \frac{a_H}{L} + (n-1) \frac{kL}{a_H} - \frac{n w_{n+1}^2 Q^2 L}{8 a_H^{2n-1}} \right], \end{aligned} \quad (9)$$

whereas the entropy S of the CFT is given by the Bekenstein-Hawking entropy (2) of the black hole without re-scaling. In terms of the energy density $\rho = E/V$, the pressure $p = \rho/n$, the charge density $\rho_Q = Q/V$ and the electrostatic potential $\Phi = \phi \frac{a}{L}$ of the CFT within the volume $V = a^n \text{Vol}(M^n)$, the Friedmann equations (6,7) take the following forms:

$$H^2 = \frac{16\pi G}{n(n-1)} \left(\rho - \frac{1}{2} \Phi \rho_Q \right) - \frac{k}{a^2}, \quad (10)$$

$$\dot{H} = -\frac{8\pi G}{n-1} (\rho + p - \Phi \rho_Q) + \frac{k}{a^2}, \quad (11)$$

where $G = (n-1)G_{n+2}/L$ is the Newton's constant on the brane. So, the cosmological evolution is determined by the energy density ρ and the pressure p of the CFT plus those of the electric potential energy due to the R-charge. From these Friedmann equations, we obtain the following energy conservation equation:

$$d(\rho - \frac{1}{2} \Phi \rho_Q)/d\tau + n(\rho + p - \Phi \rho_Q) = 0. \quad (12)$$

The Friedmann equations (10,11) can be respectively put into the following forms resembling thermodynamic formulas of the CFT:

$$S_H = \frac{2\pi}{n} a \sqrt{E_{BH} [2(E - \frac{1}{2} \Phi Q) - k E_{BH}]}, \quad (13)$$

$$k E_{BH} = n(E + pV - \Phi Q - T_H S_H), \quad (14)$$

in terms of the Hubble entropy S_H and the Bekenstein-Hawking energy E_{BH} , where

$$S_H \equiv (n-1) \frac{H V}{4G}, \quad E_{BH} \equiv n(n-1) \frac{V}{8\pi G a^2}, \quad T_H \equiv -\frac{\dot{H}}{2\pi H}. \quad (15)$$

The first Friedmann equation (10) can be expressed also as the following relation among the Bekenstein entropy $S_B \equiv \frac{2\pi a}{n} E$, the Bekenstein-Hawking entropy $S_{BH} \equiv (n-1) \frac{V}{4Ga}$, the Hubble entropy S_H and $S_Q \equiv \frac{2\pi a}{n} \cdot \frac{1}{2} \Phi Q$:

$$S_H^2 = 2(S_B - S_Q) S_{BH} - k S_{BH}^2. \quad (16)$$

For the $k = 1$ case, this can be written as the following quadratic relation:

$$S_H^2 + (S_B - S_Q - S_{BH})^2 = (S_B - S_Q)^2, \quad (17)$$

which generalizes the one in Ref. [8]. However, $S_B - S_Q$ does not remain constant during the cosmological evolution, as can be checked by applying Eq. (12) along with the equation of state $p = \rho/n$. So, one cannot draw a circular diagram with the constant radius $S_B - S_Q$ to study the time evolution of S_H and S_{BH} unlike the case in Ref. [8].

We study thermodynamics of the CFT at the moment when the brane crosses the black hole horizon $a = a_H$, defined as the largest root of $h(a) = 0$, i.e.,

$$\frac{a_H^2}{L^2} + k - \frac{w_{n+1}M}{a_H^{n-1}} + \frac{nw_{n+1}^2Q^2}{8(n-1)a_H^{2n-2}} = 0. \quad (18)$$

From Eqs. (6,18), we see that

$$H^2 = \frac{1}{L^2} \quad \text{at} \quad a = a_H. \quad (19)$$

The total entropy S of the CFT remains constant, but the entropy density,

$$s \equiv \frac{S}{V} = (n-1) \frac{a_H^n}{4GLa^n}, \quad (20)$$

varies with time. Making use of Eq. (19), we see that s at $a = a_H$ can be expressed in terms of H in the following form:

$$s = (n-1) \frac{H}{4G} \quad \text{at} \quad a = a_H, \quad (21)$$

which implies

$$S = S_H \quad \text{at} \quad a = a_H. \quad (22)$$

Making use of the general formula $H^2 = \sigma^2 - h(a)/a^2$ that follows from Eqs. (3,5), we see that the CFT temperature $T = h'(a_H)L/(4\pi a_H)$ at $a = a_H$ can be expressed in terms of H and $\dot{H}$ in the following way:

$$T = -\frac{\dot{H}}{2\pi H} \quad \text{at} \quad a = a_H. \quad (23)$$

Eq. (14) along with Eqs. (22,23) implies

$$E_C = kE_{BH} \quad \text{at} \quad a = a_H, \quad (24)$$

where E_C is the Casimir energy defined as

$$E_C \equiv n(E + pV - \Phi_H Q - TS). \quad (25)$$

So, even for the case of charged AdS black holes, for any values of k , thermodynamic quantities of the CFT take the forms simply expressed in terms of the Hubble parameter and its time derivative, when the brane crosses the black hole horizon.

Thermodynamic quantities of the CFT satisfy the first law of thermodynamics,

$$TdS = dE - \Phi_H dQ + pdV, \quad (26)$$

which can be re-expressed in terms of the densities as

$$Tds = d\rho - \Phi_H d\rho_Q + n(\rho + p - \Phi_H \rho_Q - Ts) \frac{da}{a}, \quad (27)$$

making use of $dV = nV da/a$. The combination $\rho + p - \Phi_H \rho_Q - Ts$ measures the subextensive contribution to the thermodynamic system. To find the expression for the combination, we write the energy density of the CFT in the following way:

$$\rho = \frac{na_H^n}{16\pi G_{n+2}a^{n+1}} \left(\frac{a_H}{L} + k \frac{L}{a_H} + \frac{nw_{n+1}^2 Q^2 L}{8(n-1)a_H^{2n-1}} \right), \quad (28)$$

and make use of the equation of state $p = \rho/n$, which holds for CFTs. The resulting expression is

$$\frac{n}{2}(\rho + p - \Phi_H \rho_Q - Ts) = k \frac{\gamma}{a^2}, \quad (29)$$

where the Casimir quantity γ is given by

$$\gamma = \frac{n(n-1)a_H^{n-1}}{16\pi G a^{n-1}}. \quad (30)$$

So, the Casimir energy (25) of the CFT is

$$E_C = \frac{kn(n-1)a_H^{n-1} \text{Vol}(M^n)}{8\pi G a}, \quad (31)$$

which does not have explicit dependence on Q . The entropy density (20) of the CFT can be expressed in terms of γ and the densities as

$$s^2 = \left(\frac{4\pi}{n} \right)^2 \gamma \left(\rho - \frac{1}{2} \Phi_H \rho_Q - k \frac{\gamma}{a^2} \right). \quad (32)$$

By making use of Eq. (21), we can show that the entropy density formula (32) reproduces the first Friedmann equation (10) when the brane crosses the black hole horizon. Also, making use of Eqs. (21,23), we can show that Eq. (29) reproduces the second Friedmann equation (11) when $a = a_H$. We have thus shown that the results of Ref. [9] can be extended to the charged topological AdS black hole cases.

From the entropy density formula (32), we obtain the following generalized Cardy-Verlinde formula for the CFT with the R-charge Q :

$$S = \sqrt{\frac{2\pi a}{n}} S_C [2(E - \frac{1}{2} \Phi_H Q) - E_C], \quad (33)$$

where the Casimir entropy S_C and the Casimir energy E_C are defined as

$$\begin{aligned} S_C &\equiv (n-1) \frac{a_H^{n-1} \text{Vol}(M^n)}{4G}, \\ E_C &\equiv \frac{kn}{2\pi a} S_C = kn(n-1) \frac{a_H^{n-1} \text{Vol}(M^n)}{8\pi G a}. \end{aligned} \quad (34)$$

Note, the Casimir energy E_C in the above generalized Cardy-Verlinde formula is positive, zero and negative for $k = 1, 0, -1$, respectively, whereas the Casimir entropy S_C is always positive for any k . The Casimir entropy depends implicitly on M , Q and k through a_H satisfying Eq. (18). The generalized Cardy-Verlinde formula (33) coincides with the cosmological Cardy formula (13) when the brane crosses the black hole horizon $a = a_H$. The generalized Cardy-Verlinde formula (33) can be written also as the following relation:

$$S^2 = 2(S_B - S_{Q_H})S_C - kS_C^2, \quad (35)$$

where $S_{Q_H} \equiv \frac{2\pi a}{n} \cdot \frac{1}{2} \Phi_H Q$. This relation and the relation (16) among the cosmological entropy bounds coincide when $a = a_H$.

We now discuss the cosmological holographic bounds generalized to the case of the charged topological AdS black holes. The above expressions for the generalized Cardy-Verlinde formula suggest that the criterion for distinguishing between a weakly and a strongly self-gravitating universes should be modified to

$$\begin{aligned} E - \frac{1}{2} \Phi_H Q \leq E_{BH} &\Leftrightarrow S_B - S_{Q_H} \leq S_{BH} & \text{for} & \quad Ha \leq \sqrt{2-k} \\ E - \frac{1}{2} \Phi_H Q \geq E_{BH} &\Leftrightarrow S_B - S_{Q_H} \geq S_{BH} & \text{for} & \quad Ha \geq \sqrt{2-k}. \end{aligned} \quad (36)$$

We further propose that the cosmological bound on S_C conjectured in Ref. [8] continues to hold without modification, even when $Q \neq 0$:

$$S_C \leq S_{BH}, \quad (37)$$

which can be inferred from the fact that the expression for the Casimir entropy in Eq. (34) does not explicitly depend on Q , as well as on k . From the explicit expression (34) for S_C and the definition for S_{BH} , we see that the cosmological bound (37) implies $a_H \leq a$. So, the cosmological bound (37) is saturated when $a = a_H$, i.e., when the brane crosses the black hole horizon, at which moment the entropy S , the temperature T and the Casimir energy E_C of the CFT are given respectively by Eqs. (22,23,24). We argue that the conjectured cosmological bound (37) implies the Hubble entropy bound and the modified Bekenstein bound for a strongly and a weakly self-gravitating cases, respectively. For the strongly self-gravitating case, we see from Eqs. (36,37) that $S_C \leq S_{BH} \leq S_B - S_{Q_H}$. From Eq. (35) we see that S is a monotonically increasing function of S_C in the interval $S_C \leq S_B - S_{Q_H}$ for any values of k , as long as $S_B > S_{Q_H}$. Thus, S reaches its maximum value when $S_C = S_B - S_{Q_H}$ and therefore $S_C = S_{BH}$,

for which $S = S_H$ (as can be seen from Eqs. (16,35) with $S_C = S_{BH}$ and $a = a_H$). The conjectured cosmological bound (37) therefore implies the Hubble entropy bound even for the $Q \neq 0$ case:

$$S \leq S_H \quad \text{for} \quad Ha \geq \sqrt{2-k}. \quad (38)$$

For the weakly self-gravitating case, we have $S_C \leq S_B - S_{Q_H} \leq S_{BH}$ for $k = 1$, if it is assumed that $E_C \leq E - \frac{1}{2}\Phi_H Q$. So, we have the following modified Bekenstein bound:

$$S \leq S_B - S_{Q_H} \quad \text{for} \quad Ha \leq 1, \quad (39)$$

which is saturated when $S_C = S_B - S_{Q_H}$. For any values of k , if we assume that $S_C \leq S_B - S_{Q_H}$ (which can be interpreted as the holographic upper limit on the degrees of freedom of the dual CFT, generalized to the case with an R-charge), then we also have $S_C \leq S_B - S_{Q_H} \leq S_{BH}$ for any k . The Bekenstein bound is then generalized to

$$S \leq \sqrt{2-k}(S_B - S_{Q_H}) \quad \text{for} \quad Ha \leq \sqrt{2-k}, \quad (40)$$

which is saturated when $S_C = S_B - S_{Q_H}$.

References

- [1] R. Easther and D.A. Lowe, ‘‘Holography, cosmology and the second law of thermodynamics,’’ *Phys. Rev. Lett.* **82** (1999) 4967, hep-th/9902088.
- [2] G. Veneziano, ‘‘Pre-bangian origin of our entropy and time arrow,’’ *Phys. Lett.* **B454** (1999) 22, hep-th/9902126.
- [3] D. Bak and S. Rey, ‘‘Cosmic holography,’’ *Class. Quant. Grav.* **17** (2000) L83, hep-th/9902173.
- [4] R. Brustein, ‘‘The generalized second law of thermodynamics in cosmology,’’ *Phys. Rev. Lett.* **84** (2000) 2072, gr-qc/9904061.
- [5] N. Kaloper and A. Linde, ‘‘Cosmology vs. holography,’’ *Phys. Rev.* **D60** (1999) 103509, hep-th/9904120.
- [6] R. Bousso, ‘‘A covariant entropy conjecture,’’ *JHEP* **9907** (1999) 004, hep-th/9905177; ‘‘Holography in general space-times,’’ *JHEP* **9906** (1999) 028, hep-th/9906022; ‘‘The holographic principle for general backgrounds,’’ *Class. Quant. Grav.* **17** (2000) 997, hep-th/9911002;
- [7] W. Fischler and L. Susskind, ‘‘Holography and cosmology,’’ hep-th/9806039.
- [8] E. Verlinde, ‘‘On the holographic principle in a radiation dominated universe,’’ hep-th/0008140.
- [9] I. Savonije and E. Verlinde, ‘‘CFT and entropy on the brane,’’ hep-th/0102042.
- [10] J.D. Bekenstein, ‘‘A universal upper bound on the entropy to energy ratio for bounded systems,’’ *Phys. Rev.* **D23** (1981) 287.

- [11] R. Brustein and G. Veneziano, “A causal entropy bound,” *Phys. Rev. Lett.* **84** (2000) 5695, hep-th/9912055.
- [12] R. Brustein, S. Foffa and G. Veneziano, “CFT, holography, and causal entropy bound,” *Phys. Lett.* **B507** (2001) 270, hep-th/0101083.
- [13] J.L. Cardy, “Operator content of two-dimensional conformally invariant Theories,” *Nucl. Phys.* **B270** (1986) 186.
- [14] B. Wang, E. Abdalla and R. Su, “Relating Friedmann equation to Cardy formula in universes with cosmological constant,” *Phys. Lett.* **B503** (2001) 394, hep-th/0101073.
- [15] D. Klemm, A.C. Petkou and G. Siopsis, “Entropy bounds, monotonicity properties and scaling in CFTs,” *Nucl. Phys.* **B601** (2001) 380, hep-th/0101076.
- [16] R. Cai, “The Cardy-Verlinde formula and AdS black holes,” hep-th/0102113.
- [17] A.K. Biswas and S. Mukherji, “Holography and stiff-matter on the brane,” *JHEP* **0103** (2001) 046, hep-th/0102138.
- [18] D. Birmingham and S. Mokhtari, “The Cardy-Verlinde formula and Taub-Bolt-AdS spacetimes,” hep-th/0103108.
- [19] D. Klemm, A.C. Petkou, G. Siopsis and D. Zanon, “Universality and a generalized C-function in CFTs with AdS duals,” hep-th/0104141.
- [20] D. Youm, “The Cardy-Verlinde formula and asymptotically flat charged black holes,” hep-th/0105036.
- [21] R. Cai, Y.S. Myung, and N. Ohta, “Bekenstein bound, holography and brane cosmology in charged black hole background,” hep-th/0105070.
- [22] S. Nojiri, S.D. Odintsov and S. Ogushi, “Holographic entropy and brane FRW-dynamics from AdS black hole in d5 higher derivative gravity,” hep-th/0105117.
- [23] R. Cai and Y. Zhang, “Holography and brane cosmology in domain wall backgrounds,” hep-th/0105214.
- [24] S. Nojiri and S.D. Odintsov, “AdS/CFT correspondence, conformal anomaly and quantum corrected entropy bounds,” hep-th/0011115.
- [25] S. Nojiri and S.D. Odintsov, “AdS/CFT and quantum-corrected brane entropy,” hep-th/0103078.
- [26] S. Nojiri, O. Obregon, S.D. Odintsov, H. Quevedo, and M.P. Ryan, “Quantum bounds for gravitational de Sitter entropy and the Cardy-Verlinde formula,” hep-th/0105052.
- [27] D. Youm, “The Cardy-Verlinde formula and topological AdS-Schwarzschild black holes,” hep-th/0105093.
- [28] R.B. Mann, “Pair production of topological anti-de Sitter black holes,” *Class. Quant. Grav.* **14** (1997) L109, gr-qc/9607071.
- [29] R. Cai and Y. Zhang, “Black plane solutions in four dimensional spacetimes,” *Phys. Rev.* **D54** (1996) 4891, gr-qc/9609065.

- [30] D.R. Brill, J. Louko and P. Peldan, “Thermodynamics of (3+1)-dimensional black holes with toroidal or higher genus horizons,” *Phys. Rev.* **D56** (1997) 3600, gr-qc/9705012.
- [31] R.B. Mann, “Charged topological black hole pair creation,” *Nucl. Phys.* **B516** (1998) 357, hep-th/9705223.
- [32] R. Cai, J. Ji and K. Soh, “Topological dilaton black holes,” *Phys. Rev.* **D57** (1998) 6547, gr-qc/9708063.
- [33] R. Cai and K. Soh, “Topological black holes in the dimensionally continued gravity,” *Phys. Rev.* **D59** (1999) 044013, gr-qc/9808067.
- [34] A. Chamblin, R. Emparan, C.V. Johnson and R.C. Myers, “Charged AdS black holes and catastrophic holography,” *Phys. Rev.* **D60** (1999) 064018, hep-th/9902170.
- [35] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, “Gauge theory correlators from non-critical string theory,” *Phys. Lett.* **B428** (1998) 105, hep-th/9802109.
- [36] E. Witten, “Anti-de Sitter space and holography,” *Adv. Theor. Math. Phys.* **2** (1998) 253, hep-th/9802150.
- [37] E. Witten, “Anti-de Sitter space, thermal phase transition, and confinement in gauge theories,” *Adv. Theor. Math. Phys.* **2** (1998) 505, hep-th/9803131.